

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets




(11) Publication number:

0 497 415 A1

(12)

EUROPEAN PATENT APPLICATION(21) Application number: **92200199.5**(51) Int. Cl.⁵: **C07D 453/02, A61K 31/435**(22) Date of filing: **24.01.92**(30) Priority: **01.02.91 EP 91200196**(43) Date of publication of application:
05.08.92 Bulletin 92/32(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IT LI LU MC
NL PT SE**(71) Applicant: **AKZO N.V.**
Velperweg 76
NL-6824 BM Arnhem(NL)(72) Inventor: **Wieringa, Johannes Hubertus**
Bachlaan 47
NL-5384 BL Heesch(NL)
Inventor: **van Aelst, Sjoerd Folkert**
Torenstraat 17
NL-5366 BJ Megen(NL)
Inventor: **Plate, Ralf**
Kortenhorststraat 1
NL-5344 KD Oss(NL)(74) Representative: **Hermans, Franciscus G.M. et al**
P.O. Box 20
NL-5340 BH Oss(NL)(54) **3-Quinuclidine derivatives.**

(57) The invention relates to a 3-quinuclidine derivative having the formula I

**I**

wherein X is O or S, and R represents one to five substituents independently selected from hydrogen, hydroxy, lower alkyl, lower alkoxy, lower aralkyloxy, lower alkanoyloxy, halogen, NO₂, CF₃, CN, NR₁R₂, and COR₃, in which R₁ and R₂ are independently selected from hydrogen, lower alkyl, and lower alkanoyl, and R₃ is selected from OH, lower alkyl, lower alkoxy, and NR₄R₅, wherein R₄ and R₅ are independently hydrogen or lower alkyl; or pharmaceutically acceptable salts thereof.

EP 0 497 415 A1

The invention relates to 3-quinuclidine derivatives, a process for the preparation thereof, a pharmaceutical composition containing the same, as well as to the use of these 3-quinuclidine derivatives for the preparation of a medicament.

Related quinuclidine derivatives are known, for instance (2-quinuclidinyl)-phenylether which is disclosed by C.A. Grob and A. Sieber in *Helv. Chim. Acta* 50 (8), 2531-42 (1967). This compound was obtained by fragmentation of (2-quinuclidinyl)-phenylketoxime tosylate, however, no pharmacological activity was assessed or suggested.

The present invention relates to a 3-quinuclidine derivative having the formula I



wherein X is O or S, and R represents one to five substituents independently selected from hydrogen, hydroxy, lower alkyl, lower alkoxy, lower aralkyloxy, lower alkanoyloxy, halogen, NO₂, CF₃, CN, NR₁R₂, and COR₃, in which R₁ and R₂ are independently selected from hydrogen, lower alkyl, and lower alkanoyl, and R₃ is selected from OH, lower alkyl, lower alkoxy, and NR₄R₅, wherein R₄ and R₅ are independently hydrogen or lower alkyl; or pharmaceutically acceptable salts thereof.

The compounds of this invention have muscarinic properties. They bind to muscarinic agonist receptor sites with a 2 to 750 fold preference as compared to muscarinic antagonist receptor sites, as is exemplified in their ability to bind preferentially to the agonist site of muscarinic receptors in membrane preparations of rat cerebral cortex, or membrane from rat forebrain. Preferred compounds show an agonist/antagonist binding ratio of between 10 and 400. Compounds having ratios very much higher than 750 are undesired, because they may exert side-effects or toxic properties. Compounds having similar pharmacological properties are known, e.g. pilocarpine and oxotremorine, but the chemical structures of these compounds bear no relation to the 3-quinuclidine derivatives of this invention. The 3-quinuclidine derivatives are suitable for the treatment of cognition disorders, like presenile and senile dementia, including Alzheimer's disease, and for the treatment of other cholinergic deficiencies, like Huntington's chorea, tardive dyskinesia, hyperkinesia, mania, and Tourette syndrome.

Preferred compounds according to the invention are the 3-quinuclidine derivatives in which X is O, and more preferably 3-quinuclidine derivatives in which X is O and R is selected from CF₃, bromine, or iodine. Preferably R represents one substituent in the ortho position.

The term halogen used in the definition of formula I means fluorine, chlorine, bromine, or iodine. Bromine and iodine are preferred halogens.

The term lower alkyl means a branched or unbranched alkyl group with 1-4 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, or *tert*-butyl.

The alkyl moiety which is present in the lower alkoxy group has the same meaning as previously defined for lower alkyl.

The term lower alkanoyl means an alkanoyl (acyl) group, the alkyl moiety of which has the same meaning as lower alkyl. Acetyl is the preferred lower alkanoyl group. The alkanoyl group in the lower alkanoyloxy group has the same meaning.

The term lower aralkyloxy means an aralkyloxy group, the alkyl moiety of which has 1-4 carbon atoms, and preferably is a methylene group, and the aryl moiety of which is preferably a phenyl group, optionally substituted with hydroxy, lower alkyl, lower alkoxy, or halogen. The preferred aralkyloxy group is the phenylmethoxy (benzyloxy) group.

The novel compounds of formula I may be isolated from the reaction mixture in the form of a pharmaceutically acceptable salt. The pharmaceutically acceptable salts may also be obtained by treating the free base of formula I with an organic or inorganic acid such as HCl, HBr, HI, H₂SO₄, H₃PO₄, acetic acid, propionic acid, glycolic acid, maleic acid, malonic acid, methane-sulphonic acid, fumaric acid, succinic acid, tartaric acid, citric acid, benzoic acid, and ascorbic acid.

The compounds of this invention possess a chiral carbon atom, and may therefore be obtained as a pure enantiomer, or as a mixture of enantiomers, among which the racemic mixture. Methods for obtaining the pure enantiomers are well known in the art, e.g. synthesis from chirally pure 3-quinuclidinol or 3-thioquinuclidinol, crystallization of salts which are obtained from optically active acids and the racemic mixture, or chromatography using chiral columns.

The 3-quinuclidine derivatives of the invention can be prepared by methods known for the preparation


of analogous compounds.

A suitable method is the condensation of 3-quinuclidinol or 3-thioquinuclidinol with a benzene derivative having the formula II



II

wherein R has the previously given meaning and Hal is a halogen (fluorine, chlorine, bromine, or iodine), or with a iodonium derivative having the formula III



III

wherein R has the previously given meaning, after which the compound obtained may be separated into its enantiomers and/or may be converted into a pharmaceutically acceptable salt.

It is possible to convert the products obtained by the previously mentioned procedure into another product according to the invention. Using generally known methods it is, for instance, possible to convert aromatic substituents into other aromatic substituents. Alkoxy substituents may be treated with strong acids such as BBR_3 , to give the hydroxy substituent. Hydroxy substituted compounds may be condensed with lower alcohols in acidic medium to give alkoxy derivatives, or may be acylated with acyl anhydrides to give alkanoyl substituted derivatives, and nitro substituted compounds may be reduced to afford amine compounds. Compounds wherein R_1 , R_2 , R_3 and/or R_4 are hydrogen may be alkylated, e.g. by a Leuckart-Wallach reaction, to afford compounds wherein R_1 , R_2 , R_3 and/or R_4 are alkyl. This is further exemplified in the examples.

The compounds of the invention may be administered enterally or parenterally, and for humans preferably in a daily dosage of 0,001-10 mg per kg body weight. Mixed with pharmaceutically suitable auxiliaries, e.g. as described in the standard reference, Chase et al., Remington's Pharmaceutical Sciences, the compounds may be compressed into solid dosage units, such as pills, tablets, or be processed into capsules or suppositories. By means of pharmaceutically suitable liquids the compounds can also be applied as an injection preparation in the form of a solution, suspension, emulsion, or as a spray, e.g. a nasal spray.

The invention is further illustrated by the following examples.

Example 1

3-[2-(trifluoromethyl)phenoxy]-1-azabicyclo[2.2.2]octane ethanedioate

A solution of 1.78 g (14 mmol) of 3-quinuclidinol in 20 ml of dry N,N-dimethylformamide (DMF) was added dropwise under nitrogen to a suspension of 0.6 g (15 mmol) of sodium hydride in 20 ml of dry DMF. The mixture was stirred for 15 min at 70°C , and a solution of 3.8 ml (30 mmol) of 2-fluoro-(trifluoromethyl)-benzene in 8 ml of dry DMF was added dropwise. The mixture was stirred for another 1.5 h, after which the solvent was evaporated. The residue was dissolved in ethyl acetate and washed with brine. The ethyl acetate layer was separated, dried over magnesium sulfate and evaporated to dryness. The crude compound was purified by column chromatography on silica using methanol-dichloromethane (2:8) as eluent to give 3.0 g (59% yield) of 3-[2-(trifluoromethyl)phenoxy]-1-azabicyclo[2.2.2]octane, which was converted into the oxalate salt by treatment with oxalic acid and recrystallization from methanol. m.p. 174°C .

Example 2

In an analogous manner as described in Example 1 were prepared:

- 3-phenoxy-1-azabicyclo[2.2.2]octane. m.p. 49 °C.
 3-(4-nitrophenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 242 °C.
 3-(3-nitrophenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 230 °C.
 3-(2-nitrophenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 237 °C.
 5 3-[3-(trifluoromethyl)phenoxy]-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 206 °C.
 3-[4-(trifluoromethyl)phenoxy]-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 181 °C.
 3-(2-chlorophenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 184 °C.
 3-(3-chlorophenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 196 °C.
 3-(4-chlorophenoxy)-1-azabicyclo[2.2.2]octane. m.p. 66 °C.
 10 3-(4-chlorophenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 198 °C.
 3-(3,4-dichlorophenoxy)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 129 °C.
 3-(2,6-dichlorophenoxy)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 137 °C.
 3-(2,3-dichlorophenoxy)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 179 °C.
 3-(2,4-dichlorophenoxy)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 154 °C.
 15 3-(3-fluorophenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 206 °C.
 3-(2-fluorophenoxy)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 92 °C.
 3-(2-fluorophenoxy)-1-azabicyclo[2.2.2]octane ethanedioate. m.p. 174 °C.
 3-(2-bromophenoxy)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 139 °C.
 3-(2-iodophenoxy)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 159 °C.
 20 3-(4-methoxyphenoxy)-1-azabicyclo[2.2.2]octane (oil).
 3-(3-methoxyphenoxy)-1-azabicyclo[2.2.2]octane (oil).
 3-(2-methoxyphenoxy)-1-azabicyclo[2.2.2]octane hydrochloride. m.p. 166 °C.
 3-[2-(phenylmethoxy)phenoxy]-1-azabicyclo[2.2.2]octane (oil).
 2-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]phenol (Z)-2-butenedioate (obtained by catalytic hydrogenation of 3-[2-
 25 (phenylmethoxy)phenoxy]-1-azabicyclo[2.2.2]octane). m.p. 126 °C.
 2-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]phenol acetate (Z)-2-butenedioate (obtained by acetic anhydride/K₂CO₃
 treatment of 2-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]phenol). m.p. 161 °C.
 2-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenamine dihydrochloride (obtained by catalytic hydrogenation of 3-
 (2-nitrophenoxy)-1-azabicyclo[2.2.2]octane). m.p. >250 °C.
 30 3-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenamine dihydrochloride (obtained by catalytic hydrogenation of 3-
 (3-nitrophenoxy)-1-azabicyclo[2.2.2]octane).
 4-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenamine dihydrochloride (obtained by catalytic hydrogenation of 3-
 (4-nitrophenoxy)-1-azabicyclo[2.2.2]octane). m.p. >250 °C.
 N-[4-(1-azabicyclo[2.2.2]oct-3-yl)oxy]phenylacetamide hydrochloride (obtained by acetylation of 4-[(1-
 35 azabicyclo[2.2.2]oct-3-yl)oxy]benzenamine). m.p. >250 °C.
 2-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzonitrile (Z)-2-butenedioate. m.p. 137 °C.
 3-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzonitrile (Z)-2-butenedioate.
 4-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzonitrile (Z)-2-butenedioate.
 2-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenecarboxamide (obtained by KOH/t-butanol hydrolysis of 2-[(1-
 40 azabicyclo[2.2.2]oct-3-yl)oxy]benzonitrile). m.p. 144 °C.
 4-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenecarboxamide hydrochloride. m.p. >250 °C.
 3-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenecarboxamide hydrochloride. m.p. >250 °C.
 2-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzoic acid.

45
 Example 3

4-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenamine dihydrochloride.

- 50 3-(4-nitrophenoxy)-1-azabicyclo[2.2.2]octane (Example 2) was submitted to catalytic hydrogenation with
 Pd/C in ethanol, after which the free base was converted into the hydrochloric salt to obtain 4-[(1-
 azabicyclo[2.2.2]oct-3-yl)oxy]benzenamine dihydrochloride. M.p. >250 °C.

Example 4

- 55 N-[4-(1-azabicyclo[2.2.2]oct-3-yl)oxy]phenylacetamide hydrochloride.

4-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]benzenamine (Example 3) was acetylated with acetic anhydride in

dry pyridine at -10 °C. The crude product was converted into the hydrochloric salt to obtain N-[4-[(1-azabicyclo[2.2.2]oct-3-yl)oxy]phenyl]acetamide hydrochloride. M.p. >250 °C.

Example 5

In an analogous manner as described in Example 1 were prepared from (S)- or (R)-3-quinuclidinol: (S)-3-(2-fluorophenoxy)-1-azabicyclo[2.2.2]octane ethanedioate. m.p. 158 °C; $[\alpha]_D^{20} = +27.5^\circ$ (c = 1, water).

(R)-3-(2-fluorophenoxy)-1-azabicyclo[2.2.2]octane ethanedioate. m.p. 159 °C; $[\alpha]_D^{20} = -27.5^\circ$ (c = 1, water).

Example 6

In an analogous manner as described in Example 1, using 3-thioquinuclidinol (K.B. Shaw, Can. J. Chem., 43 (1965) 3112) instead of 3-quinuclidinol were prepared:

3-(2-nitrophenylthio)-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 156 °C.

3-[2-(trifluoromethyl)phenylthio]-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 120 °C.


3-[4-(trifluoromethyl)phenylthio]-1-azabicyclo[2.2.2]octane (Z)-2-butenedioate. m.p. 127 °C.

2-[(1-azabicyclo[2.2.2]oct-3-yl)thio]benzenamine (Z)-2-butenedioate (obtained by iron/hydrochloric acid reduction of 3-(2-nitrophenylthio)-1-azabicyclo[2.2.2]octane). m.p. 106 °C.

N-[4-[(1-azabicyclo[2.2.2]oct-3-yl)thio]phenyl]acetamide (Z)-2-butenedioate.

Claims


1. A 3-quinuclidine derivative having the formula I



I

wherein X is O or S, and R represents one to five substituents independently selected from hydrogen, hydroxy, lower alkyl, lower alkoxy, lower aralkyloxy, lower alkanoyloxy, halogen, NO₂, CF₃, CN, NR₁R₂, and COR₃, in which R₁ and R₂ are independently selected from hydrogen, lower alkyl, and lower alkanoyl, and R₃ is selected from OH, lower alkyl, lower alkoxy, and NR₄R₅, wherein R₄ and R₅ are independently hydrogen or lower alkyl; or pharmaceutically acceptable salts thereof.

2. The 3-quinuclidine derivative of claim 1, wherein X is O; or pharmaceutically acceptable salts thereof.
3. The 3-quinuclidine derivative of claim 1 or 2, wherein R is selected from CF₃, bromine and iodine; or pharmaceutically acceptable salts thereof.
4. The 3-quinuclidine derivative of any of claims 1-3, wherein R represents a substituent in the ortho position.
5. The 3-quinuclidine derivative of any of claims 1-4 for use in therapy.
6. A process for the preparation of the 3-quinuclidine derivative of any of claims 1-4, characterized in that 3-quinuclidinol or 3-thioquinuclidinol is condensed with a benzene derivative having the formula II




II

wherein R has the previously given meaning and Hal is a halogen, or with a iodonium derivative having

the formula III

5



III

10

wherein R has the previously given meaning, after which the compound obtained may be separated into its enantiomers and/or may be converted into a pharmaceutically acceptable salt.

15

7. A pharmaceutical composition comprising the 3-quinuclidine derivative of any of claims 1-4 in admixture with pharmaceutically acceptable auxiliaries.
8. A use of the 3-quinuclidine derivative of any of claims 1-4, for the manufacture of a medicament for the treatment of cognition disorders and cholinergic deficiencies.

20

25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 20 0199
PAGE1

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	EP-A-0 370 415 (BOEHRINGER INGELHEIM KG) * page 3, line 41 - line 46; claims 1,4 * ---	1,5,7,8	C07D453/02 A61K31/435
A	EP-A-0 190 920 (A.H. ROBINS CO., INC.) * claims 1,7 * ---	1,5,7,8	
A	EP-A-0 158 532 (A.H. ROBINS CO., INC.) * page 1, line 2 - line 8; claims 1,2 * ---	1,5,7	
A	EP-A-0 311 724 (DELALANDE S.A. / A.H. ROBINS CO., INC.) * claims 1,5-7 * ---	1,5,7	
A	EP-A-0 221 702 (BEECHAM GROUP PLC) 1,10,12 ---	1,5,7	
A	GB-A-2 208 385 (J. WYETH AND BROTHER LTD.) * abstract; claims 1-3 * ---	1,5,7	
A	CHEMICAL ABSTRACTS, vol. 99, no. 23, 5 December 1983, Columbus, Ohio, US; abstract no. 187523G, E.R. ATKINSON ET AL.: 'Parasympatholytic ...' page 78 ; & J. Med. Chem. 1983, 26(12), 1772-5 ---	1,5,7	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
A	CHEMICAL ABSTRACTS, vol. 86, no. 21, 23 May 1977, Columbus, Ohio, US; abstract no. 155489R, E. MIKHLINA ET AL.: 'Synthesis and ...' page 456 ; Khim.-Farm. Zh. 1976, 10(11), 56-60 ---	1,5,7	C070
A,D	HELVETICA CHIMICA ACTA, vol. 50, no. 8, 11 December 1967, pages 2531 - 2542; C.A. GROB ET AL.: 'Fragmentierung von ...' * page 2534, compound 21 * -----	1	
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 13 MAY 1992	Examiner HASS C.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document			

